

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number  
**WO 03/093222 A1**

(51) International Patent Classification<sup>7</sup>: C07C 233/63,  
A61K 31/198

(21) International Application Number: PCT/IN02/00114

(22) International Filing Date: 29 April 2002 (29.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): BIOCON  
INDIA LIMITED [IN/IN]; 20th K.M. Hosur Road, Heb-  
bagodi, Bangalore 561 229, Karnataka (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RAJAMAHEN-  
DRA, Shanmughasamy [IN/IN]; Biocon India Limited,  
20th Km. Hosur Road, Hebbagodi, Bangalore 561 229,  
Karnataka (IN). ASWATHANARAYANAPPA, Chan-  
drashekar [IN/IN]; Biocon India limited, 20th Km Hosur  
Road, Hebbagodi, Bangalore 561 229, Karnataka (IN).  
PUTHIAPARAMPIL, Tom, Thomas [IN/IN]; Biocon  
India Limited, 20th Km Hosur Road, Hebbagodi, Banga-  
lore 561 229, Karnataka (IN). SRIDHARAN, Madhavan  
[IN/IN]; Biocon India Limited, 20th Km Hosur Road, Heb-  
bagodi, Bangalore 561 229, Karnataka (IN). GANESH,

Sambasivam [IN/IN]; Biocon India Limited, 20th Km  
Hosur Road, Hebbagodi, Bangalore 561 229, Karnataka  
(IN).

(74) Agents: ANAND, Pravin et al.; Anand & Anand Advo-  
cates, B-41, Nizamuddin East, New Delhi 110 013 (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

[Continued on next page]

(54) Title: NOVEL FORM OF N-(TRANS-4-ISOPROPYLCYCLOHEXYLCARBONYL)-D-PHENYLALANINE



(57) Abstract: Novel polymorph Form C of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine is produced having different infra red spectrum and X-ray diffraction patterns from previously known forms of the compound.

WO 03/093222 A1



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Novel Form Of**  
**N- (trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine**

**FIELD OF THE INVENTION**

5

The present invention relates to a novel crystalline form of N- (trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine and a process for preparing the same.

10 **BACKGROUND OF THE INVENTION**

N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine has therapeutic utility in depressing blood glucose levels in the management of Type 2 diabetes mellitus.

15

N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine is disclosed in Japanese Patent Application No. 63-54321 (equivalent to EP-A-196222 and US 4,816,484). The Japanese application describes how the compound may be crystallized from aqueous methanol to yield crystals having a melting point of 20 129°C to 130° C. These crystals are referred as "B-type". These B-type crystals suffer from problems of instability, especially when subjected to mechanical grinding.

J. Med. Chem. 32, 1436 (1989) discusses preparation of N-  
25 (cyclohexylcarbonyl)-D-phenylalanines and related compounds which includes N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine.

US Patent Nos. 5,463,116 and 5,488,150 discloses preparation of novel crystalline form of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine designated "H-type". This patent also discloses a method of treating form "B" crystals with appropriate solvent mixtures to obtain form "H" crystals and vice-versa. The H-type crystals have an enhanced stability over B type crystals.

Yaowu Fenxi Zazhi 2001, 21, 342 discusses "S" form of nateglinide which is different from form "B" and form "H".

## 10 SUMMARY OF THE INVENTION

According to a first aspect of the invention there is provided a novel process for the production of novel crystalline form "C" of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine by reacting D-phenylalanine methylester HCl with *trans*-4-isopropylcyclohexane carboxylic acid in presence of propane phosphonic acid anhydride or LiOH-Al<sub>2</sub>O<sub>3</sub> in halogenated hydrocarbon solvents such as dichloromethane, dichloroethane at a temperature between -10° C to 90° C followed by base hydrolysis.

20 Alternatively, the product can be obtained by reacting *trans*-4-isopropylcyclohexane carbonyl chloride with D-phenylalanine methylester HCl in halogenated hydrocarbon solvents such as dichloromethane, dichloroethane in presence of base such as triethylamine, pyridine at a temperature between -10° C to 90° C followed by base hydrolysis.

25

According to a still further aspect of the present invention, the new crystal from "C" of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine thus produced have at least one, and preferably all, of the following properties:

- 5 (a) a melting point in the range of 126 °C to 132 °C;
- (b) a powder X-ray diffraction pattern comprising characteristic peaks at 14.0, 17.8, 19.0, 20.2 and  $21.2 \pm 0.2$  degrees measured at reflection angle  $2\theta$ ;
- (c) an infrared absorption spectrum comprising absorption bands at with  
10 absorption bands in the region of 1742, 1648, 1599, 1540 and  $1191 \pm 2$   
cm<sup>-1</sup>.

Yet another aspect of the invention includes a N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine Form C for use in the  
15 preparation of a medicament for treating Type 2 diabetes mellitus.

This invention also includes a pharmaceutical composition comprising a therapeutically effective amount of crystals of Form C N-(trans-4-isopropylcyclohexylcarbonyl)-D-Phenylalanine.

20

Another aspect of the invention, relates to a method for treating patients suffering from type 2 diabetes mellitus by administering a therapeutically effective amount of the pharmaceutical composition of Form C of N-(trans-4-isopropylcyclohexylcarbonyl)-D-Phenylalanine.

25

## BRIEF DESCRIPTION OF THE ACCOMPANYING DRAWINGS

FIG. 1 shows a powder X-ray diffraction pattern of form C crystals of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine;

FIG. 2 shows an infra red absorption spectrum of form C crystals of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine;

## DETAILED DESCRIPTION OF THE INVENTION

The process according to the inventors for preparing N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine form C comprises

- (a) suspending N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester in water or a water miscible solvent;
- (b) treating the suspension with a base;
- (c) addition of water followed by adjusting the pH to 1.0-4.0 using a mineral acid;
- (d) extracting using ethyl acetate;
- (e) concentrating the ethyl acetate extract;
- (f) addition of petroleum ether to the ethyl acetate concentrate; and
- (g) filtering and drying the precipitate obtained to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine form C.

The water miscible solvent in step (a) is selected from methanol, ethanol, isopropanol, or a mixture of these, preferably methanol. The base in step (b) is selected from potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide or a mixture of these preferably potassium carbonate.

The precipitate obtained in step (g) is suspended in water before filtration and drying to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine form C.

5 N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester is prepared by:

- reacting D-Phenylalanine methyl ester hydrochloride with trans-4-isopropylcyclohexane carboxylic acid in the halogenated hydrocarbons
- 10 - filtering the reaction mixture;
- concentrating to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.

The first step in the above process for the preparation of N-(trans-4-  
15 isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester is carried out in the presence of propane phosphonic acid anhydride, LiOH adsorbed in aluminium oxide or triethylamine.

An alternate method for producing N-(trans-4-  
20 isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester comprises the steps of

- reacting D-Phenylalanine methyl ester hydrochloride with trans-4-isopropylcyclohexane carbonyl chloride in halogenated hydrocarbon solvents and in the presence of a base,
- 25 - filtering the reaction mixture;
- concentrating to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.



The halogenated hydrocarbon solvents are selected from dichloromethane or dichloroethane.

- 5 The base are selected from triethylamine or pyridine. The the reaction temperature is -10 ° to 90 ° C.

Embodiments of the invention are illustrated below by way of the following examples, not to be considered as limiting.

10

#### EXAMPLE 1

N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester:

- D-Phenylalanine methyl ester hydrochloride (10g, 0.046 mol) was suspended in a solution of triethylamine (33 mL) in dichloromethane (50 mL); the mixture  
15 was cooled to 0-5 °C and trans-4-isopropylcyclohexane carboxylic acid (7.9 g, 0.046 mol) was added to the reaction mixture. A solution of Propane phosphonic acid anhydride (46.4 mL, 0.046 mol) in ethyl acetate was added dropwise over a period of 30 minutes, maintaining the temperature at 0-5 °C and stirred for 14 hours at ambient temperature. The reaction mixture was washed with 1.5 N HCl,  
20 5% sodium bicarbonate solution and brine. The organic layer was concentrated to yield 12.5 g of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.

#### EXAMPLE 2

- 25 N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester:

D-Phenylalanine methyl ester hydrochloride (20g, 0.092 mol) was suspended in a solution of triethylamine (66 mL) in dichloroethane (100 mL) and the mixture was stirred for 1 h at room temperature. The organic layer was separated after washing with water and dried over anhydrous sodium sulphate. trans-4-  
5 Isopropylcyclohexane carboxylic acid (15.8 g, 0.092 mol) was added to the organic layer followed by LiOH-Al<sub>2</sub>O<sub>3</sub> (5.5g of LiOH adsorbed on 40.5g Aluminum oxide) and refluxed for 24 hours. The reaction mixture was filtered over celite bed and washed with 1.5 N HCl, 5% sodium bicarbonate solution and brine. The organic layer was concentrated to yield 15.5 g of N-(trans-4-  
10 isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.

### EXAMPLE 3

N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester:

D-Phenylalanine methyl ester hydrochloride (26g, 0.12 mol) was suspended in a  
15 solution of triethylamine (85 mL) in dichloromethane (125 mL) and the mixture was cooled to 0-5 °C. A solution of trans-4-isopropylcyclohexane carbonyl chloride (25g, 0.13 mol) in dichloromethane (75 mL) was added dropwise over a period of 10 minutes, maintaining the temperature at 0-5° C and stirred for 12 hours at ambient temperature. The reaction mixture was washed with 1.5 N HCl,  
20 5% sodium bicarbonate solution and brine. The organic layer was concentrated to yield 38g of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.

### EXAMPLE 4

25

N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine:

To suspension of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester (38 g, 0.11 mol) in methanol (600 mL), a solution of potassium carbonate (80g, 0.57 mol) in water (400 mL) was added and the reaction mixture was stirred for 12 h at ambient temperature. Water (1500 mL) was added and pH  
5 was adjusted to 2.0 by adding 6N HCl. The mixture was extracted using ethyl acetate (3 x 400 mL) and the combined extract was washed with brine. The organic layer was concentrated to about 150 mL and pet. ether (300 mL) was added. The product was filtered and suspended in water (600 mL) and stirred for 12 hours at ambient temperature. The slurry  
10 was filtered and dried to yield 35g of the title compound.

The compound showed a sharp melting point of 128 – 129 °C. X-ray diffraction pattern and infra red absorption spectrum of the final compound was recorded and identified as form C crystals of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine.

15

**We claim:**

1. Crystalline N-(trans-4-isopropylcyclohexylcarbonyl)-D-Phenylalanine  
5 having at least one of the following properties:
  - (a) a melting point in the range of 126° to 132 ° C;
  - (b) a powder X-ray diffraction pattern comprising characteristic  
peaks at 14.0, 17.8, 19.0, 20.2 and  $21.2 \pm 0.2$  degrees  
measured at reflection angle  $2\theta$ ;
  - 10 (c) an infrared absorption spectrum comprising absorption bands  
at with absorption bands in the region of 1742, 1648, 1599,  
1540 and  $1191 \pm 2 \text{ cm}^{-1}$ .
2. N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine Form C.  
15
3. N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine Form C for use  
in the preparation of a medicament for treating Type 2 diabetes mellitus.
4. A process for preparing N-(trans-4-isopropylcyclohexylcarbonyl)-D-  
20 phenylalanine form C comprising the steps of:
  - (d) suspending N-(trans-4-isopropylcyclohexylcarbonyl)-D-  
phenylalanine methyl ester in water or a water miscible  
solvent;
  - (e) treating the suspension with a base;
  - 25 (f) addition of water followed by adjusting the pH to 1.0-4.0  
using a mineral acid;
  - (d) extracting using ethyl acetate;

- (e) concentrating the ethyl acetate extract;
- (f) addition of petroleum ether to the ethyl acetate concentrate;
- (g) filtering and drying the precipitate obtained to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine form C.

5

5. A process of claim 4 wherein the water miscible solvent in step (a) is selected from methanol, ethanol, isopropanol, or a mixture of these.
6. A process of claim 5 wherein the water miscible solvent in step (a) is  
10 methanol.
7. A process of claim 4 wherein the base in step (b) is selected from potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, lithium hydroxide or a mixture of these.  
15
8. A process of claim 7 wherein the base in step (b) is potassium carbonate.
9. A process of claim 4-8 wherein the precipitate obtained in step (g) is  
20 suspended in water before filtration and drying to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine form C.
10. A process of claim 4 wherein N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester of step (a) is prepared by:  
25
  - reacting D-Phenylalanine methyl ester hydrochloride with trans-4-isopropylcyclohexane carboxylic acid in the halogenated hydrocarbons

- filtering the reaction mixture;
- concentrating to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.

- 5 11. A process of claim 10 wherein said halogenated hydrocarbon solvents are selected from dichloromethane or dichloroethane.
12. A process of claim 10 wherein step (a) is carried out in the presence of propane phosphonic acid anhydride, LiOH adsorbed in  
10 aluminium oxide or triethylamine.
13. A process of claim 4 wherein N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester of step (a) is prepared by:
- reacting D-Phenylalanine methyl ester hydrochloride  
15 with trans-4-isopropylcyclohexane carbonyl chloride in halogenated hydrocarbon solvents and in the presence of a base,
  - filtering the reaction mixture;
  - concentrating to get N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine methyl ester.  
20
14. A process of claim 13 wherein said halogenated hydrocarbon solvents are selected from dichloromethane or dichloroethane.
- 25 15. A process of claim 13 wherein said base are selected from triethylamine or pyridine.

16. A process of claim 10 or 13 wherein the reaction temperature is  
-10 ° to 90 ° C.
17. A pharmaceutical composition comprising a therapeutically  
s effective amount of crystals of Form C N-(trans-4-  
isopropylcyclohexylcarbonyl)-D-Phenylalanine according to claim 1.

FIG 1







FIG 2



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/IN02/00114

## A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl. <sup>7</sup>: C07C 233/63; A61K 31/198

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN: Chemical Abstracts; based on the compound of the invention in the claims

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 196 222 B1 (AJINOMOTO CO., INC.) 1 October 1986 See example 31, (m.p.). (& US 4,816,484 A)	1-17
X	Shinkai H et al.: "N-(Cyclohexylcarbonyl)-D-phenylalanines and Related Compounds. A New Class of Oral Hypoglycemic Agents. 2"; Journal of Medicinal Chemistry, 1989, Vol. 32 NO. 7, pp 1436-41. See table II, compound 13	1-17
A	US 5,463,116 A (Sumikawa et al.) 31 October 1995 Whole document	1-17



Further documents are listed in the continuation of Box C



See patent family annex

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

5 September 2002

Date of mailing of the international search report

17 SEP 2002

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE  
PO BOX 200, WODEN ACT 2606, AUSTRALIA  
E-mail address: pct@ipaustalia.gov.au  
Facsimile No. (02) 6285 3929

Authorised officer

G. D. HEARDER

Telephone No : (02) 6283 2553

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/IN02/00114

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstracts 136:159110; Li G et al.: "A new crystal structure in nateglinide found by X-ray powder diffraction"; Yaowu Fenxi Zazhi (2001), 21(5), 342-344 Abstract	1-17

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/IN02/00114

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member			
EP	196222	US	4816484	JP	63054321
US	5463116	BR	1100807	CA	2114678
		LU	90843	LU	90846
		JP	5208943	EP	526171
				US	5488150
END OF ANNEX					